

Very large set axioms over Constructive set theories

Hanul Jeon

Cornell University

2022-04-29

Cornell Logic Seminar

This work is joint work with Richard Matthews (Leeds)

Table of Contents

- 1 Review: Large cardinals
- 2 Constructive set theories
- 3 Main results

Measurable cardinals

- Defined in terms of ultrafilters
- Scott (1961) associated measurable cardinals with elementary embeddings:

Definition

Let $M \subseteq V$ be a transitive class.

- $j: V \rightarrow M$ is an elementary embedding if j respects all first-order formulas, that is,

$$V \models \phi(\vec{a}) \iff M \models \phi(j(\vec{a})).$$

- The critical point $\text{crit } j$ of an elementary embedding $j: V \rightarrow M$ is the least ordinal moved by j .


Theorem (Scott 1961)

A cardinal κ is measurable if and only if it is a critical point of some elementary embedding $j: V \rightarrow M$.

Theorem (Scott 1961)

Measurable cardinals do not exist in L .

Generalizations of measurable cardinals were extensively studied by various people. (For example, Solovay-Reinhardt-Kanamori 1978).



Reinhardt's dream

The attempt to find a stronger notion of large cardinal bore the notion now known as a Reinhardt cardinal.

Definition

A Reinhardt cardinal is a critical point of an elementary embedding $j: V \rightarrow V$.

Flying too close to the sun – Kunen Inconsistency

Kunen (1971) proved that Reinhardt's notion cannot be realized over ZFC:

Theorem (Kunen Inconsistency theorem, 1971)

Work over ZFC, there is no Reinhardt cardinals.

In fact, there is no elementary embedding $j: V_{\lambda+2} \rightarrow V_{\lambda+2}$

It is still open whether a Reinhardt cardinal is compatible with ZF, that is, in the choiceless context.

Night Flying: Choiceless large cardinals

Some difficulties in working without the axiom of choice:

- Not all cardinal correspond to an ordinal.
- Not all successor cardinal are regular.
- Constructions and proofs become harder.
- Needs to consider more before applying the known results.

Super Reinhardt cardinals

Definition

A cardinal κ is

- Super Reinhardt if for each ordinal α we can find an elementary embedding $j: V \rightarrow V$ such that $\text{crit } j = \kappa$ and $j(\kappa) > \alpha$.
- A-super Reinhardt for a class A if for each ordinal α we can find an elementary embedding for A-formulas $j: V \rightarrow V$ such that $\text{crit } j = \kappa$ and $j(\kappa) > \alpha$.

Super Reinhardt sets were defined by Woodin in 1983, and extensively studied in the mid-2010s in the context of Woodin's HOD-dichotomy.

Totally Reinhardt cardinals

Definition

- We call Ord is totally Reinhardt if for every class A we can find a cardinal κ which is A -super Reinhardt.
- κ is totally Reinhardt if $(V_\kappa, V_{\kappa+1}) \models \text{Ord is totally Reinhardt}$.

The Wholeness axiom

Before defining the Wholeness axiom, let us analyze the definition of a Reinhardt cardinal:

Definition

ZFC with a Reinhardt cardinal is a theory comprising:

- Language: \in and a unary function symbol j ,
- Axioms: Usual axioms of ZFC, with the elementarity of j , and Separation and Replacement for j -formulas.

Definition (Corazza 2000)


The Wholeness Axiom WA is obtained by restricting Replacement to formulas with no j .

We can further weaken WA as follows:

Definition

The theory Basic Theory of Elementary Embedding (BTEE) is claim that $j: V \rightarrow V$ is a elementary embedding. BTEE does not subsume Separation and Replacement for j -formulas.

That is, we obtain BTEE by dropping Separation for j -formulas from WA.



IZF and CZF: A brief history

- (H. Friedman, 1973) Intuitionistic ZF (IZF) with the double-negation translation between IZF and ZF.
- Various attempts to formalize the foundation for Bishop-styled constructive mathematics.
- Myhill's constructive set theory CST.
- (Aczel, 1978) Constructive ZF and its type-theoretic interpretation.

Axioms of ZF

Definition

- 1 Extensionality: $a = b \iff \forall x(x \in a \leftrightarrow x \in b)$.
- 2 Pairing: $\{a, b\}$ exists.
- 3 Union: $\bigcup a$ exists.
- 4 Separation: $\{x \in a \mid \phi(x)\}$ exists,
- 5 Replacement: $\{F(x) \mid x \in a\}$ exists if F is a class function.
- 6 Power set: $\mathcal{P}(a) = \{x \mid x \subseteq a\}$ exists.
- 7 Regularity: Every set has a \in -minimal element.
- 8 Infinity: \mathbb{N} exists.

Axioms of IZF

Definition

- 1 Extensionality: $a = b \iff \forall x(x \in a \leftrightarrow x \in b)$.
- 2 Pairing: $\{a, b\}$ exists.
- 3 Union: $\bigcup a$ exists.
- 4 Separation: $\{x \in a \mid \phi(x)\}$ exists.
- 5 **Collection**: if $\forall x \in a \exists y \phi(x, y)$, then there is b such that $\forall x \in a \exists y \in b \phi(x, y)$
- 6 Power set: $\mathcal{P}(a) = \{x \mid x \subseteq a\}$ exists.
- 7 **Set Induction**: $\forall a[[\forall x \in a \phi(x)] \rightarrow \phi(a)] \rightarrow \forall a \phi(a)$
- 8 Infinity: \mathbb{N} exists.

Axioms of CZF

Definition

- 1 Extensionality: $a = b \iff \forall x(x \in a \leftrightarrow x \in b)$.
- 2 Pairing: $\{a, b\}$ exists.
- 3 Union: $\bigcup a$ exists.
- 4 **Bounded Separation**: $\{x \in a \mid \phi(x)\}$ exists if ϕ is bounded.
- 5 **Strong Collection**: if $\forall x \in a \exists y \phi(x, y)$, then there is b such that $\forall x \in a \exists y \in b \phi(x, y)$ and $\forall y \in b \exists x \in a \phi(x, y)$.
- 6 **Subset Collection**: There is a full subset of $\text{mv}(a, b)$.
- 7 **Set Induction**: $\forall a [[\forall x \in a \phi(x)] \rightarrow \phi(a)] \rightarrow \forall a \phi(a)$
- 8 Infinity: \mathbb{N} exists.

Differences between IZF and CZF



IZF

- 1 Full separation
- 2 Powerset
- 3 Impredicative
- 4 Equiconsistent with ZF

and more...



CZF

- 1 Bounded separation
- 2 Subset collection
- 3 Allows type-theoretic interpretation
- 4 Far more weaker than ZF

Large set axioms

- Ordinals over constructive set theories are not well-behaved.

Examples (CZF)

- Every ordinal is well-ordered \implies Excluded Middle for Δ_0 -formulas,
 - $\alpha \subseteq \beta$ does not imply $\alpha \in \beta$ or $\alpha = \beta$.
-
- We define large cardinal properties over constructive set theories by mimicking the structural properties of H_κ and V_κ .

Multi-valued functions

The notion of multi-valued function provides a syntactic sugar for Collection.

Definition (Multi-valued function)

Let A and B be classes. We call a relation R of domain A and codomain B a multi-valued function from A to B . (Notation: $R: A \rightrightarrows B$)

Definition (Subimage)

If $R: A \rightrightarrows B$ and $R^{-1}: B \rightrightarrows A$, we write $R: A \Leftrightarrow B$, and we call B a subimage of R .

Multi-valued functions replace functions in CZF-context.

From Replacement to Strong Collection

Definition (Replacement)

If $F: a \rightarrow V$ is a first-order definable class function, then we can find a set image b of F .

Definition (Strong Collection)

If $R: a \rightrightarrows V$ is a first-order definable class **multi-valued** function, then we can find a set **subimage** b of R . (That is, $R: a \Leftrightarrow b$.)

From Powerset to Subset Collection

Definition (Powerset, equivalent formulation)

For given sets a, b , we can find a set c such that...

If $f: a \rightarrow b$, then c contains an image of f . (i.e., $\text{Im } f \in c$.)

Definition (Subset Collection, equivalent formulation)

For given sets a, b , we can find a set c such that...

If $r: a \rightrightarrows b$, then c contains a **subimage** of r . (i.e., there is $d \in c$ such that $r: a \rightrightarrows d$.)

Defining large sets

With some pain, we can prove

Lemma (ZFC, regular cardinal)

A cardinal κ is regular if and only if for $a \in H_\kappa$ and $f: a \rightarrow H_\kappa$, we have $\text{Im } f \in H_\kappa$.

By mimicking the above result, we have

Definition (CZF, regular set)

A transitive set K is regular if $a \in K$, $r \in K$ and $r: a \rightrightarrows K$, then we can find some subimage $b \in K$ of r (that is, $r: a \rightleftarrows b$.)

Similarly, we can see

Lemma (ZFC, inaccessible cardinal)

A cardinal κ is inaccessible if and only if κ is regular and H_κ satisfies:


- 1** $\omega \in H_\kappa$, H_κ is closed under union and intersection, and
- 2** if $a, b \in H_\kappa$, then $c := \{\text{Im } f \mid f: a \rightarrow b\} \in H_\kappa$.

Definition (CZF, inaccessible set)

A set K is inaccessible if K is regular and

- 1** $\omega \in K$, K is closed under union and intersection, and
- 2** if $a, b \in K$, then we can find $c \in K$ such that we can always find a subimage of $r: a \rightrightarrows b$, $r \in K$ from c .

Consistency hierarchy



Large sets and elementary embeddings

Let us consider elementary embeddings over CZF:

Definition

Let $j: V \rightarrow M$ be an elementary embedding. A set K is a critical point of j if K is the 'least' set lifted by j in the sense that



- $j(x) = x$ for all $x \in K$, and
- $K \in j(K)$

Definition

A set K is critical if K is inaccessible and a critical point of an elementary embedding $j: V \rightarrow M$.

Question: the consistency strength of CZF with a critical cardinal.

Two scenarios



A Lower bound

Theorem (J., Matthews, CZF)

Let K be a critical point of a Σ_0 -elementary embedding $j: V \rightarrow M$ such that K satisfies Δ_0 -separation. Then $K \models \text{IZF}$.

(Note: the above theorem does not require Separation, Strong Collection or Set Induction for j -formulas.)

Theorem

CZF with a critical set proves the consistency of ZFC + BTEE

Reinhardt embeddings

Definition

An inaccessible set K is a Reinhardt set if K is a critical point of $j: V \rightarrow V$.

Theorem

CZF with a Reinhardt cardinal proves $\text{Con}(\text{ZF} + \text{WA})$.

Go beyond the Reinhardtness

Definition

- An inaccessible set K is super Reinhardt if for every set a we can find an elementary embedding $j: V \rightarrow V$ such that K is a critical point of j and $a \in j(K)$.
- An inaccessible set K is A -super Reinhardt if for every set a we can find an A -elementary embedding $j: V \rightarrow V$ such that K is a critical point of j and $a \in j(K)$.

Theorem

CZF with a super Reinhardt set proves the consistency of ZF with a Reinhardt cardinal.

Definition


V is totally Reinhardt (abbr. V is TR) is the following claim: for every class A , there is an A -super Reinhardt set K .

Theorem

CZF with 'V is TR' proves all axioms of IZF. Furthermore, the following two theories are equiconsistent:

- $\text{CZF} + \text{'V is TR,}'$ and
- $\text{ZF} + \text{'V is TR.}'$

(The exact definition for super/totally Reinhardt require the formulation of constructive second-order set theory [Appendix](#))



A rough sketch for the proofs

The proof divides into 2-3 main steps:

- 1** Internal analysis of the given large set axioms. Usually produces a model of $IZF + X$.
- 2** Double-negation translation: Friedman-styled translation, Gambino's Heyting-valued model, or their combinations. The resulting lower bound is of the form $\text{Con}(ZF + X)$
- 3** If possible, derive the consistency strength in terms of ZFC with large cardinal axioms.

Open problems

- 1 Non-trivial upper bounds for the consistency strength.
- 2 Better lower bounds. (For example, can we derive $\text{Con}(\text{ZFC} + \text{WA})$ from $\text{Con}(\text{ZF} + \text{WA})$?)
- 3 Defining other large set notions (e.g., supercompactness and extendibles) and analyzing their consistency strength.
- 4 Questions regarding machinery in the paper, e.g., second-order constructive set theory.

Questions



Thank you!

Constructive second-order set theory

Definition (Constructive Gödel-Bernays set theory, CGB)

CGB is defined over the two-sorted languages (sets and classes) with the following axioms:

- Axioms of CZF for sets.
- Every set is a class, and every element of a class is a set.
- Class Extensionality: two classes are equal if they have the same set members.
- Elementary Comprehension: if $\phi(x, p, C)$ is a first-order formula with a class parameter C , then there is a class A such that $A = \{x \mid \phi(x, p, C)\}$.

Definition (CGB, Continued)

- Class Set Induction:

$$\forall^1 A [[\forall^0 x (\forall^0 y \in x (y \in A) \rightarrow x \in A)] \rightarrow \forall^0 x (x \in A)].$$

- Class Strong Collection:

$$\forall^1 R \forall^0 a [R: a \rightrightarrows V \rightarrow \exists^0 b (R: a \leftrightarrow b)].$$

Definition (Intuitionistic Gödel-Bernays set theory, IGB)

IGB is obtained by adding the following axioms to CGB:

- Axioms of IZF for sets.
- Class Separation: if A is a class and a is a set, then $A \cap a$ is a set.

Note that CGB and IGB are conservative extensions of CZF and IZF respectively.

The definition of an elementary embedding $j: V \rightarrow M$ requires quantifying over formulas ϕ :

$$\phi(\vec{a}) \iff \phi^M(j(\vec{a})).$$

We resolve this problem by introducing the infinite conjunction \bigwedge .

Definition (CGB with the infinite connectives, CGB_∞)

CGB_∞ has the same axiom with CGB, but defined over the first-order intuitionistic logic with the infinite connectives \bigwedge and \bigvee .

Super Reinhardt sets and 'V is TR' are defined over CGB_∞ . Also, CGB_∞ is a conservative extension of CGB.

(Back to [main](#).)